

The Efficiency of Transfer-matrix Simulations of Supramolecular Magnets in the Parallel Computing Environment

Ryszard Matysiak and Grzegorz Kamieniarz

*Institute of Engineering and Computer Education, University of Zielona Góra
ul. prof. Z. Szafrana 4, 65-516 Zielona Góra, Poland
r.matysiak@eti.uz.zgora.pl*

*Computational Physics Division, Institute of Physics, Adam Mickiewicz University
ul. Umultowska 85, 61-614 Poznań, Poland
gjk@amu.edu.pl*

(Rec. 11 November 2005)

Abstract: The deterministic quantum transfer-matrix (QTM) technique and its mathematical background are presented. This important tool in computational physics can be applied to a class of the real physical low-dimensional magnetic systems described by the Heisenberg hamiltonian which includes the macroscopic molecular-based spin chains, small size magnetic clusters embedded in some supramolecules and other interesting compounds. In order to convert existing application for the susceptibility calculations to run on the grid, the speed-up and efficiency of parallelization are analyzed on the SGI Origin 3800 platform with $p = 128$ processor units. Using Message Parallel Interface (MPI) system libraries we find the efficiency of the code of 94% for $p = 128$ that makes our application suitable for the grid.

Key words: quantum transfer-matrix technique, Message Parallel Interface

1. INTRODUCTION

In this article we describe numerical simulation based on the quantum transfer-matrix (QTM) method, we present some applications in computational physics and analyze the characteristics of parallelization which make our code suitable to run on the grid. The domain of applications of the QTM method is very wide and includes not only simulations of the thermodynamic properties of the low-dimensional quantum systems [1], but also phase transitions [2].

Before presenting the model and the numerical QTM technique, we review physical systems belonging to the class of low-dimensional magnets, in particular the molecular magnets. First we consider the compound $[\text{Mn}(\text{hfac})_2\text{NITPh}]_6$ (hfac, hexafluoroacetyl acetone; NITPh, 2-phenyl-4,4,5,5-tetramethyl-4,5-dihydro-1H-imidazolyl-1-oxy-3-oxide) belonging to a class of nano-compounds actively investigated for their magnetic properties [3, 4]. Synthesis of polynuclear metal complexes with oxygen atom bridges has resulted in a series of new molecules with unusual geometric symmetries and patterns [5]. Their magnetic properties, associated to a large number of interacting paramagnetic centers in a single aggregate, have significantly stimulated the research effort with the prospect of technological applications [3]. The interest in spin assemblies stems from the fact that they set the low-size limit for

magnetic nanoparticles. They can display magnetic quantum tunneling [5] and quantum-size effects in the thermodynamical properties [6].

2. MODEL AND SIMULATION TECHNIQUE

To simulate the finite-temperature properties of low-dimensional magnetic systems, we model rings or chains in the framework of the spin Hamiltonian with the nearest neighbor interaction, which can be described by the following operator:

$$\mathcal{H} = -J \sum_{i=1}^N (\mathbf{S}_i \cdot \mathbf{S}_{i+1}) - D \sum_{i=1}^N (S_i^z)^2 - g_v \mu_B B \sum_{i=1}^N S_i^v, \quad (1)$$

where \mathbf{S}_i is interpreted as the spin located at the i -th site of a one-dimensional lattice of N equally spaced sites. N may become infinite for the macroscopic chain. J denotes the nearest neighbor exchange integral (negative for the anti-ferromagnetic coupling) and D stands for the anisotropy parameter. B is the external magnetic field which can be applied along the chain ($v = z$) or in the perpendicular direction ($v = x, y$), g_v is the corresponding gyromagnetic ratio and N is the size of a given one-dimensional system (the chain or the ring). The spin values S_i may be uniform or non-uniform and define the matrix representation of the corresponding non-commuting operators.

We calculate the thermodynamic properties from the derivatives of the free energy related to the partition function Z . For the spin system described in (1) we can calculate the canonical partition function Z from the definition

$$Z = \text{Tr} e^{-\beta \mathcal{H}}$$

The Hilbert space of states of an N -site low-dimensional system is a direct product of single spin spaces, therefore, the base states can be labelled by the N -tuple of the eigenvalues of the z component of the single spin operator

$$|S_1^z \dots S_N^z\rangle \equiv |S_1^z\rangle \dots |S_N^z\rangle. \quad (3)$$

The values of matrix elements of $e^{-\beta \mathcal{H}}$ cannot be calculated for large N because of non-commuting operators in (1). Thus, to eliminate this restriction, we look for systematic approximants to the partition function Z .

We express Hamiltonian (1) in terms of the spin-pair operators $\mathcal{H}_{i, i+1}$ as

$$\begin{aligned} \mathcal{H} = -\sum_{i=1}^N & \left\{ J \mathbf{S}_i \cdot \mathbf{S}_{i+1} + \frac{1}{2} D \left[(S_i^z)^2 + (S_{i+1}^z)^2 \right] + \right. \\ & \left. + \frac{1}{2} g_\nu \mu_B B (S_i^\nu + S_{i+1}^\nu) \right\}. \end{aligned} \quad (4)$$

In the checker-board decomposition (CBD) we divide the Hamiltonian (4) into two non-commuting parts [7]

$$\begin{aligned} \mathcal{H} &= \mathcal{H}^{\text{odd}} + \mathcal{H}^{\text{even}} = \\ &= (\mathcal{H}_{1,2} + \dots + \mathcal{H}_{N-1,N}) + (\mathcal{H}_{2,3} + \dots + \mathcal{H}_{N,1}), \end{aligned} \quad (5)$$

each part defined by the commuting spin-pair operators $\mathcal{H}_{i, i+1}$. Then the series of the classical approximants of the quantum thermal values can be found, using the general Suzuki-Trotter formula [7]. The partition function is calculated from the expression

$$\mathcal{Z} = \lim_{m \rightarrow \infty} \mathcal{Z}_m = \lim_{m \rightarrow \infty} \text{Tr} \left[\prod_{i=1}^{N/2} v_{2i-1, 2i} \prod_{i=1}^{N/2} v_{2i, 2i+1} \right]^m, \quad (6)$$

where $v_{i, i+1} = e^{-\beta \mathcal{H}_{i, i+1}/m}$, $i = 1, 2, \dots, N$ and m is a natural number (referred to as the Trotter number).

The approximant Z_m can be calculated numerically, without any restrictions on the value of N , by the quantum transfer-matrix method. The computation of Z_m is possible for relatively small values of m in the case of macroscopic chains, because of computer storage limitation, but the leading errors in taking a finite m approximant are of the order of $1/m^2$ and therefore, extrapolations to $m \rightarrow \infty$ can be performed. However, for small finite systems (e.g. Mn_6 [1]), the QTM approach can be applied for very large m , as the computing time increases linearly with m .

The trace in Eq. (6) is taken over all the configurations of the classical Ising variable S_{ir} (the eigenvalues of S_{ir}^z) on a planar lattice of the size $N \times 2m$. The lattice is obtained in the checker-board decomposition and sketched in Fig. 1. The shaded squares denote the corresponding Boltzmann weights $v_{i, i+1}$ present in (6) which are determined by some 4-spin operators.




Fig. 1. The structure of the classical map of the quantum ring of the alternating spins in the checker-board decomposition. The shaded squares denote the Boltzmann vertices, whereas the full circles and the open circles stand for spins at the odd and even sites of chain

The thermodynamic functions for all systems described by Hamiltonian (1) are related to the free energy which can be calculated from the formula $\mathcal{F} = -k_B T \ln Z$. The specific heat is given as its second derivative with respect to temperature, the magnetization is then evaluated from the first and the zero-field susceptibility from the second derivative with respect to the field.

3. SIMULATION RESULTS

The QTM method has been applied to simulation of the $S = 1$ one-dimensional Heisenberg model and our results are compared with the experimental results for many compounds [1]. We have studied the finite temperature static properties in wide ranges of temperatures and the single-ion anisotropy (the D parameter in Eq. (1)).

Here we report the susceptibility of the Mn_6 molecule [1]. Depending on temperature and the number n of the spin pairs $S = 1/2$ and $S = 5/2$, in some cases the calculations preserving 5 decimal places required as many as $m = 900$ steps in the Trotter direction. To save the supercomputer time, for temperatures higher than $k_B T / |J| = 0.05$ we performed our simulations only up to $n = 5$ pairs. Then we extrapolated the estimates of the zero-field susceptibility to $n = 6$ which corresponded to Mn_6 . The uncertainties, smaller than 1% at higher tem-

peratures, reached the order of 5% for about $k_B T/J = 0.05$ and at this point the QTM simulations were performed for $n = 6$ pairs.

From the computational point of view, the diagonal elements of the transfer matrix with size $12^{n=6} = 2985984$ had to be calculated. Their number was reduced approximately by a factor of 6 due to the symmetry considerations so that we had to evaluate about 0.5×10^6 matrix elements. It is a formidable task typical for the currently synthesized magnetic supramolecules [8] and needs effective methods of parallelization on the large number of processors and ultimately the grid environment.




Fig. 2. The speed-up S_p of the parallel computation for the number of processors $p \leq 128$ tested on the SGI Origin 3800 supercomputer. The line describe the function $S_p = p$

In the remaining part of our work we analyze the properties of our code with respect to the distribution of the computational task over a number of processors available in the Supercomputing and Networking Center in Poznań.

We have performed tests for a model molecule consisting of $n = 4$ pairs with $S_1 = 1/2$ and $S_2 = 5/2$ corresponding to the real Mn_6 cluster. The size of the transfer matrix is equal 20736 and a number of independent diagonal matrix elements amounts to about 5000.

The performance of our parallelized code is presented in Fig. 2. We have run the program in the SGI Origin 3800 supercomputer. To reach the convergence for our simplified model system, we need in the sequential version of the algorithm about 14600 CPU seconds for the SGI platform. The speed-up of the parallelization based on the MPI libraries is drawn by circles in Fig. 2. The speed-up has been computed as the quotient of the CPU time of the sequential version of the algorithm divided by the maximum

CPU time used by the slave processes plus the master process CPU time [9, 10].

4. CONCLUSIONS

We have worked out QTM approach to characterize the finite temperature magnetic properties of the high nuclearity cyclic spin clusters with large and alternating spins and a number of the macroscopic quasi-one-dimensional magnets.

Using large-scale computations, high resolution data for the $S = 1$ spin chains down to low temperatures and in the wide range of single-ion anisotropy parameters [1] can be found. For Mn_6 cluster the QTM technique provides the numerically exact results [1]. Our simulations demonstrate that the QTM approach is a valuable tool for calculations of the finite-temperature properties.

The computational complexity of our problems is exponential so that for the finite ring simulations we are limited by the CPU time resources and a new grid environment should be envisaged. The efficiency of the parallel computation is diminished from 100% for $p \leq 4$ to 94% for $p = 128$ but still very high and encouraging.

Acknowledgements

We thank Dr N. Meyer and Mr M. Zawadzki for many discussions and suggestions related to the parallel implementation of our application. This work was supported in part by the MNiSzW grant No. 4T11F 014 24 and the MAGMANet project NMP3-CT-2005-515767. Numerical calculations were carried out on the local computing cluster and the PSNC supercomputing platforms.

References

- [1] G. Kamieniarz and R. Matysiak, Comp. Mat. Sci. **28**, 353 (2003).
- [2] G. Kamieniarz, Phase Transitions **57**, 105 (1996).
- [3] D. Gatteschi, A. Caneschi, L. Pardi and R. Sessoli, Science **265**, 1054 (1994).
- [4] A. Caneschi, D. Gatteschi, C. Sangregorio, R. Sessoli, L. Sorace, A. Cornia, M. A. Novak, C. Paulsen and W. Wernsdorfer, J. Magn. Magn. Mat. **200**, 182 (1999).
- [5] D. Gatteschi, R. Sessoli and A. Cornia, J. Chem. Soc., Chem. Commun. 725 (2000).
- [6] S. G. Louie, Nature **384**, 612 (1996).
- [7] T. Delica and H. Leschke, Physica **A176**, 736 (1990).
- [8] H. Andres *et al.*: Chem. Eur. J. **21** 4867 (2002).
- [9] E. F. Van de Velde, Concurrent Scientific Computing, Springer-Verlag New York, Inc. (1994).
- [10] J. Błażejewicz, J. Kaczmarek, M. Kasprzak and J. Weglarz, Computational Methods in Science and Technology **1**, 31 (1996).
- [11] A. Caneschi, D. Gatteschi, J. Laugier, P. Rey, R. Sessoli, and C. Zanchini, J. Am. Chem. Soc. **110**, 2795 (1988).



RYSZARD MATYSIAK received the Ph. D. degree in Physics Science from the A. Mickiewicz University in Poznań in 1999. Currently he works in the Institute of Engineering and Computer Education University of Zielona Góra. His field of interest covers simulations of the low-dimensional magnetic systems including molecular nanomagnets.

GRZEGORZ KAMIENIARZ is a professor in theoretical and computational physics at the Institute of Physics of the Adam Mickiewicz University in Poznań. His field of interest covers statistical mechanics, phase transitions, phenomenological modeling and simulations of the low-dimensional magnetic systems including molecular nanomagnets.